AccuSpike-IR Product Specification and Analysis Report

TEL/FAX: 504-895-3338 or 866-895-3338

EMAIL: custserv@waterborneinc.com

Product: 2.0 mL, self-standing, polypropylene microtube containing 100 each, counted by flow-cytometry cell sorting, of *Giardia lamblia, H3 isolate, cysts and Cryptosporidium parvum, Iowa isolate, oocysts suspended in* 0.75 mL of a de-ionized water/0.01% Tween 20 solution. AccuSpike-IR is designed for percent recovery determination with matrix (environmental) and reagent water samples by US EPA Methods 1622, 1623, and 1623.1.

Species/genus identification method: Direct immunofluorescence microscopy with genus-specific monoclonal antibodies; also phase microscopy.

Purification method: Cysts and oocysts are purified from feces by sucrose and Percoll density gradient centrifugation.

Quantitation method: Cells enumerated using flow cytometry.

Storage Conditions: 4 C. DO NOT FREEZE.

Inactivation/Sterilization: Gamma irradiation.

Lot# 143

Expiration: 05/08/2024 for Giardia & Cryptosporidium analysis

Expiration: 06/05/2024 for Cryptosporidium analysis ONLY

Calibration Data

Calibration Data		
Test:	Spike Dreporation	Snike Dremaration
	Spike Preparation	Spike Preparation
Organism:	Cryptosporidium	Giardia
Source of Organism:	Experimentally infected mice.	Experimentally infected gerbils.
Organism strain:	CpAZ	H3
Stock suspension lot number:	20624-119	240208
•		
Date cells collected by source:	2/6/24	2/8/24
Date of initial calibration:	2/13/24	2/13/24
Age of cells (in days):	7	5
Storage media:	e-ionized Water/0.01% Tween 2	e-ionized Water/0.01% Tween 20
Storage temperature:	4 C	4 C
Viability (PI) before Irradiation	99.8%	99.5%
Mean of the counts:	100.00	99.70
Standard deviation of the coun	1.20	1.10
Relative standard deviation:	1.20	1.10

Notes from Sorting Facility:

⁽¹⁾ Mean, standard deviation and relative standard deviation are calculated from a minimum of 12 calibration verification samples per set of 10 standards.

⁽²⁾ Parasites are evaluated for general quality and intactness under DIC microscopy prior to use.